

List of Figures

Figure 1.1	North Australian locations mentioned in the text.	2
Figure 2.1	Northeast Arnhem Land region and the location of the Point Blane Peninsula study area.	16
Figure 2.2	Blue Mud Bay physiographic divisions.	17
Figure 2.3	Geological divisions for the Point Blane Peninsula and neighbouring areas.	18
Figure 2.4	Dominant gravelly, earthy sands of the Point Blane Peninsula.	19
Figure 2.5	Exposed lateritic profiles on the coastal margin of Myaoola Bay.	19
Figure 2.6	Expansive saltflats in the Grindall Bay area.	20
Figure 2.7	Estuarine meanders and dendritic tidal channels across elevated black soil plains.	20
Figure 2.8	Example of a coastal quartzite outcropping, Point Blane Peninsula.	21
Figure 2.9	(A) Global sea-level curve for the past 140 ka; (B) Stages of sea-level rise for the past 20 ka for northern Australia and surrounding region.	22
Figure 2.10	Smoothed sea-level curve for the Gulf of Carpentaria. Grey band represents the marine/non-marine transition period for Lake Carpentaria.	23
Figure 2.11	Holocene estuarine sedimentation/progradational model.	25
Figure 2.12	(A) Australian climatic regions; (B) Mean rainfall, mean temperature and mean relative humidity at Gove Airport, northern Arnhem Land.	26
Figure 2.13	The timing and nature of Holocene climatic changes, wet phases indicated by grey shading.	28
Figure 2.14	The hydrology of the Point Blane Peninsula and neighbouring areas.	30
Figure 2.15	Section of freshwater wetland towards the mouth of the Durabudboi River.	31
Figure 2.16	Durabudboi River billabong.	31
Figure 2.17	The distribution of the main vegetation regimes in the Point Blane Peninsula study area and neighbouring areas.	33
Figure 2.18	Diagrammatic representation of Yolngu seasonal calendar, northeast Arnhem Land from the data gathered by Barber (2005 90) contained in Table 2.3.	36
Figure 3.1	Survey transects locations on the Point Blane Peninsula.	42
Figure 3.2	The distribution of sites on the Point Blane Peninsula by site type.	45
Figure 3.3	Location of sites on the Point Blane Peninsula with radiocarbon determinations obtained from surface samples.	47
Figure 3.4	Calibrated radiocarbon age estimates (1 and 2 σ) from surface samples across the Point Blane Peninsula.	47
Figure 3.5	(A) Site location relative to the present and hypothetical maximum extent of sea levels; (B) Site location plotted against height above present sea level (2m contour interval) and present coastline (0m).	48
Figure 3.6	Distance to present coastline plotted against height above present sea level for (A) Myaoola Bay and (B) Grindall Bay.	50

Figure 3.7	The distribution of all sites relative to sources of freshwater.	51
Figure 3.8	The percentage of sites by minimum distance to water interval.	53
Figure 3.9	The percentage of sites containing molluscan species by the total number of sites and broad locality.	56
Figure 3.10	Density map of the number of molluscan taxa per site across the study area (contour interval = two species exploited).	57
Figure 3.11	The distribution of sites containing stone artefacts on the Point Blane Peninsula.	62
Figure 3.12	Stone artefact densities across the Point Blane Peninsula.	65
Figure 3.13	Location of quartzite and silcrete artefacts across the Point Blane Peninsula, with 2km (thin line) and 4km (thick line) intervals from quartzitic outcrop.	66
Figure 4.1	Distribution of excavated sites used in this analysis (in bold) and their Yolngu locality name (also indicated are the other excavated sites on the peninsula).	72
Figure 4.2	Location of the Yilpara community and the site of BMB/018 at Gurranganydji.	73
Figure 4.3	Site of BMB/018 at Gurranganydji, location of vehicle track relative to shell deposits, with shell eroding from the edge of the track (arrow) and the density of shell deposited on ground surface (circled).	74
Figure 4.4	BMB/018 stratigraphic diagram, south section, showing conventional radiocarbon age and 2 σ calibrated age range.	75
Figure 4.5	Density (g/cm ³) of excavated components (shell, bone, coral/pumice and laterite/rock) by excavation unit, BMB/018, 6mm sieve fraction.	76
Figure 4.6	Location of site of BMB/067b at Gurratjinya.	77
Figure 4.7	Location of site of BMB/067b Test-pit 1 at the base of a large beach-ridge, bordering a small seasonal swamp at Gurratjinya (location of excavation indicated by position of ranging pole).	78
Figure 4.8	BMB/067b Test-pit 1 stratigraphic diagram, west section, showing conventional radiocarbon ages and 2 σ calibrated age ranges.	78
Figure 4.9	BMB/067b Test-pit 1 NISP: MNI fragmentation ratios for <i>Anadara granosa</i> and <i>Marcia hiantina</i> per excavation unit, 6mm sieve fraction.	79
Figure 4.10	Density (g/cm ³) of shell, bone/otoliths, charcoal and laterite/rock by excavation unit, BMB/067b Test-pit 1, 6mm sieve fraction.	81
Figure 4.11	Lumatjpi cluster and location of site BMB/084 (excavated and radiocarbon dated site BMB/116 also shown).	82
Figure 4.12	BMB/084 site plan at Lumatjpi, showing concentrations of midden material and the location of Test-pits.	83
Figure 4.13	BMB/084 Test-pit 1 stratigraphic diagram, west section, showing conventional radiocarbon ages and 2 σ calibrated age ranges.	84
Figure 4.14	BMB/084, Test-pit 1, NISP: MNI fragmentation ratios for <i>Marcia hiantina</i> per excavation unit, 6mm sieve fraction.	85
Figure 4.15	Density (g/cm ³) of shell, charcoal, calcareous worm shell and laterite/rock by excavation unit, BMB/084 Test-pit 1, 6mm sieve fraction.	86
Figure 4.16	Garangarri cluster and location of excavated site BMB/029 (radiocarbon dated site BMB/033 also shown).	87

Figure 4.17	Site BMB/029, view west towards edge of laterite ridge and wetlands.	88
Figure 4.18	Contour plan of site BMB/029 showing location of the excavation square.	88
Figure 4.19	BMB/029 stratigraphic diagram, east section, showing conventional radiocarbon ages and 2 σ calibrated age ranges.	89
Figure 4.20	Calibrated radiocarbon ages (1 and 2 σ) obtained for BMB/029.	90
Figure 4.21	BMB/029 NISP: MNI fragmentation ratios for <i>Anadara granosa</i> and <i>Marcia hiantina</i> per excavation unit, 6mm sieve fraction.	91
Figure 4.22	Density (g/cm ³) of shell, bone/otolith, charcoal and laterite/rock by excavation unit, BMB/029, 6mm sieve fraction.	92
Figure 4.23	Dilmitjpi cluster and location of excavated site BMB/071 (excavated and radiocarbon dated site BMB/061 also shown).	93
Figure 4.24	Contour plan of site BMB/071 showing location of the excavation square.	95
Figure 4.25	Site BMB/071, view west towards the edge of the laterite ridge and dense monsoon vine thicket.	95
Figure 4.26	BMB/071 stratigraphic diagram, east section, showing conventional radiocarbon ages and 2 σ calibrated age ranges.	96
Figure 4.27	BMB/071 NISP: MNI fragmentation ratios for <i>Anadara granosa</i> and <i>Macra abbreviata</i> molluscan species per excavation unit, 6mm sieve fraction.	96
Figure 4.28	Density (g/cm ³) of shell, bone/otolith, charcoal and laterite/rock by excavation unit, BMB/071, 6mm sieve fraction.	98
Figure 4.29	Burrpilingbuy cluster and location of excavated site BMB/045 (radiocarbon dated sites BMB/036 and BMB/052 also shown).	99
Figure 4.30	Site BMB/045, view south-west towards the extensive mangrove forest.	99
Figure 4.31	Contour plan of site BMB/045 showing location of the excavated area.	100
Figure 4.32	BMB/045 stratigraphic diagram, east section, showing conventional radiocarbon ages and 2 σ calibrated age ranges.	101
Figure 4.33	BMB/045 NISP: MNI fragmentation ratio for <i>Anadara granosa</i> and <i>Macra abbreviata</i> per excavation unit, 6mm sieve fraction.	102
Figure 4.34	Density (g/cm ³) of shell, bone/otoliths, charcoal and laterite/rock by excavation unit, BMB/045, 6mm sieve fraction.	104
Figure 5.1	BMB/018, percentages of the number of taxa and volume-corrected MNI estimates (MNI/cm ³) per habitat.	114
Figure 5.2	BMB/018, dominant molluscan species by MNI percentages and volume-corrected MNI (MNI/cm ³).	115
Figure 5.3	BMB/067b, corrected MNI values (MNI/cm ³) for the dominant species by excavation unit.	118
Figure 5.4	BMB/084, corrected MNI values (MNI/cm ³) for molluscan habitat by excavation unit.	120
Figure 5.5	BMB/084, corrected MNI values for the dominant species by excavation unit.	122
Figure 5.6	BMB/029, corrected MNI values (MNI/cm ³) for molluscan habitat by excavation unit (note differences in scale).	126

Figure 5.7	BMB/029, corrected MNI values for the dominant species by excavation unit.	128
Figure 5.8	BMB/071, corrected MNI values (MNI/cm ³) for molluscan habitats by excavation unit (note differences in scale).	130
Figure 5.9	BMB/071, corrected MNI values for the dominant species by excavation unit.	131
Figure 5.10	BMB/045, volume-corrected MNI values for the dominant species by excavation unit.	134
Figure 6.1	Comparison of the 1 and 2 σ calibrated radiocarbon ages for sites on Myaoola Bay (above), and the calibrated radiocarbon age summed probability plot (below).	143
Figure 6.2	Comparison of the 1 and 2 σ calibrated radiocarbon ages for sites on Grindall Bay (above), and the calibrated radiocarbon age summed probability plot (below).	143
Figure 6.3	The distribution of site clusters (shell deposits only) on the margins of the Point Blane peninsula.	146
Figure 6.4	Growth rates for <i>Anadara granosa</i> from commercial culture beds in Malaysia.	148
Figure 6.5	<i>Anadara granosa</i> valve terminology and measured attributes for predicting maximum valve length from morphometric equations.	151
Figure 6.6	Size frequency distributions drawn from Malaysian commercial plots A, B, C, D and E; live collected sample from natural shell bed near Kokato Island, north Queensland.	153
Figure 6.7	<i>Anadara granosa</i> size frequency distributions from the Grindall Bay mounds (BMB/029, BMB/071, BMB/045) and midden (BMB/061); and from the Myaoola Bay midden BMB/067b. Black line indicates expected valve size at which sexual maturity is reached (25mm).	156
Figure 6.8	Boxplots of <i>Anadara granosa</i> valve size from the Grindall Bay mounds (BMB/029, BMB/071, BMB/045) and midden (BMB/061); and from the Myaoola Bay midden BMB/067b (dashed line indicates significant difference at 0.05 level).	160
Figure 6.9	<i>Anadara granosa</i> mean valve size / 95% confidence interval (dashed black line indicates significant difference at 0.05 level, with dashed grey line indicating possible recovery and subsequent decline); and % of juveniles (<25mm) by excavation unit / age cal BP.	162
Figure 7.1	Chronological variation in the exploitation of mollusc species by habitat, with calibrated radiocarbon age summed probability plots for Myaoola and Grindall Bays (grey shading indicates phase of significant <i>A. granosa</i> size recovery in the Grindall Bay sites).	169
Figure 7.2	Comparison of the calibrated radiocarbon summed probability distributions (BP) from <i>Anadara granosa</i> dominated mound deposits in coastal regions across northern Australia (grey shading indicates principal phase of mound formation).	171
Figure 7.3	Holocene palaeoenvironmental patterns for Australasia and the Indo-Pacific contrasted with the major period of mound formation (3000 to 500 BP) across northern Australia.	174